

University of Groningen

## Enantioselective Synthesis of 2-Aryl-4-piperidones via Rhodium/Phosphoramidite-Catalyzed Conjugate Addition of Arylboroxines

Jagt, Richard B.C.; Vries, Johannes G. de; Minnaard, Adriaan J.; Feringa, Bernard

*Published in:*  
ChemInform

*DOI:*  
[10.1002/chin.200544164](https://doi.org/10.1002/chin.200544164)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2005

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Jagt, R. B. C., Vries, J. G. D., Minnaard, A. J., & Feringa, B. (2005). Enantioselective Synthesis of 2-Aryl-4-piperidones via Rhodium/Phosphoramidite-Catalyzed Conjugate Addition of Arylboroxines. *ChemInform*, 36(44). <https://doi.org/10.1002/chin.200544164>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Enantioselective Synthesis of 2-Aryl-4-piperidones via Rhodium/Phosphoramidite-Catalyzed Conjugate Addition of Arylboroxines


Richard B. C. Jagt, Johannes G. de Vries,\* Ben L. Feringa,\* and Adriaan J. Minnaard\*

Department of Organic and Molecular Inorganic Chemistry, Stratingh Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

j.g.de.vries@rug.nl; b.l.feringa@rug.nl; a.j.minnaard@rug.nl

Received April 6, 2005

## ABSTRACT



The highly enantioselective synthesis of 2-aryl-4-piperidones by rhodium/phosphoramidite-catalyzed conjugate addition of arylboroxines to 2,3-dihydro-4-pyridones is described. Both enantiomers of a variety of products with sterically and electronically different R substituents were obtained in high isolated yield and with excellent enantioselectivity up to 99%.

The piperidine ring system is a frequently encountered heterocyclic unit in natural compounds and drug candidates.<sup>1</sup> Piperidine alkaloids exhibit a range of biological activities and as such represent important synthetic targets. Piperidones serve an important role as intermediates *en route* to substituted piperidines<sup>2</sup> and can be found as a part of more complex biologically active compounds.<sup>3</sup> Therefore, the development of short, enantioselective routes to substituted piperidones is a major goal.<sup>4</sup> An attractive catalytic route toward enantiopure piperidones is based on the enantioselective conjugate addition to readily accessible *N*-protected 2,3-dihydro-4-pyridones,<sup>5</sup> which are frequently used in alkaloid

synthesis.<sup>6</sup> Until recently,<sup>7–9</sup> however, no suitable procedures had been developed.

Monodentate phosphoramidite ligands comprise a cheap and easily tunable class of ligands that has already proven to be successful in a variety of reactions, including rhodium-catalyzed asymmetric hydrogenations,<sup>10</sup> rhodium-catalyzed conjugate additions of trifluoroborates<sup>11</sup> and boronic acids,<sup>12</sup> and copper-catalyzed asymmetric conjugate additions of

(1) For a general review on piperidines, see: Rubiralta, M.; Giralt, E.; Diez, A. *Piperidine: Structure, Preparation and Synthetic Applications of Piperidine and its Derivatives*; Elsevier: Amsterdam, 1991.

(2) For some recent examples of the use of 4-piperidones in the synthesis of biologically active 4-piperidines, see: (a) Tawara, J. N.; Lorenz, P.; Stermitz, F. R. *J. Nat. Prod.* **1999**, *62*, 321–323. (b) Watson, P. S.; Jiang, B.; Scott, B. *Org. Lett.* **2000**, *2*, 3679–3681. (c) Brooks, C. A.; Comins, D. L. *Tetrahedron Lett.* **2000**, *41*, 3551–3553.

(3) For an example, see: Mukhtar, T. A.; Wright, G. D. *Chem. Rev.* **2005**, *105*, 529–542.


(4) (a) For a review of the recent developments in asymmetric routes to substituted piperidines, see: Bailey, P. D.; Millwood, P. A.; Smith, P. D. *Chem. Commun.* **1998**, 633–640. (b) For a review on the recent advances in the synthesis of piperidines and piperidones, see: Weintraub, P. M.; Sabol, J. S.; Kane, J. M.; Borcharding, D. R. *Tetrahedron* **2003**, *59*, 2953–2989.

(5) For the preparation of *N*-protected 2,3-dihydropyridones, see: (a) Kozikowski, A. P.; Park, P.-U. *J. Org. Chem.* **1990**, *55*, 4668–4682. (b) Comins, D. L.; Chung, G.; Foley, M. A. *Heterocycles* **1994**, *37*, 1121–1140.

(6) For a review on the asymmetric preparation and synthetic utility of 2,3-dihydro-4-pyridones, see: Comins, D. L. *J. Heterocycl. Chem.* **1999**, *36*, 1491–1500.

(7) Shintani, R.; Tokunaga, N.; Doi, H.; Hayashi, T. *J. Am. Chem. Soc.* **2004**, *126*, 6240–6241.

(8) Šebesta, R.; Pizzuti, M. G.; Boersma, A. J.; Minnaard, A. J.; Feringa, B. L. *Chem. Commun.* **2005**, 1711–1713.



**Figure 1.** Phosphoramidite **L**, a highly efficient ligand in the rhodium-catalyzed conjugate addition of boronic acids.

diorganozinc reagents.<sup>13</sup> Recently, we have reported the synthesis of 2-alkyl-4-piperidones with high enantiomeric excess using the copper/phosphoramidite-catalyzed conjugate addition of dialkylzinc reagents to *N*-protected 2,3-dihydro-4-pyridones.<sup>8</sup> It was noted that this type of substrate is less reactive toward 1,4-addition than cyclic enones, *e.g.*, 2-cyclohexenone.

Although highly enantioselective 1,4-addition of diphenylzinc, using the same catalyst, has been reported for 2-cyclohexenone,<sup>14</sup> the lack of readily available diarylzinc reagents severely limits this method. A more convenient method for the introduction of aryl and alkenyl moieties is the asymmetric rhodium-catalyzed conjugate addition of boronic acids pioneered by Hayashi and Miyaura.<sup>15</sup> Excellent levels of enantioselectivity have been achieved for a broad range of enones using BINAP.<sup>15</sup> Also phosphonites<sup>16</sup> and amidophosphines<sup>17</sup> were successfully applied as chiral ligands. It was shown by our group that phosphoramidites (*i.e.*, **L**, Figure 1) are exceptionally efficient ligands for this reaction in terms of reaction rate, chemoselectivity, and enantioselectivity.<sup>12</sup>

(9) For related 1,4-additions to 2-piperidones, see: Pineschi, M.; Del Moro, F.; Gini, F.; Minnaard, A. J.; Feringa, B. L. *Chem. Commun.* **2004**, 1244–1245.

(10) (a) Van den Berg, M.; Minnaard, A. J.; Schudde, E. P.; Van Esch, J.; de Vries, A. H. M.; de Vries, J. G.; Feringa, B. L. *J. Am. Chem. Soc.* **2000**, *122*, 11539–11540. (b) Peña, D.; Minnaard, A. J.; de Vries, J. G.; Feringa, B. L. *J. Am. Chem. Soc.* **2002**, *124*, 14552–12553. (c) Van den Berg, M.; Minnaard, A. J.; Haak, R. M.; Leeman, M.; Schudde, E. P.; Meetsma, A.; Feringa, B. L.; de Vries, A. H. M.; Maljaars, C. E. P.; Willans, C. E.; Hyett, D.; Boogers, J. A. F.; Henderickx, H. J. W.; de Vries, J. G. *Adv. Synth. Catal.* **2003**, *345*, 308–323.

(11) (a) Duursma, A.; Boiteau, J.-G.; Lefort, L.; Boogers, J. A. F.; de Vries, A. H. M.; de Vries, J. G.; Minnaard, A. J.; Feringa, B. L. *J. Org. Chem.* **2004**, *69*, 8045–8052. (b) Duursma, A.; Lefort, L.; Boogers, J. A. F.; de Vries, A. H. M.; de Vries, J. G.; Minnaard, A. J.; Feringa, B. L. *Org. Biomol. Chem.* **2004**, *2*, 1682–1684.

(12) (a) Boiteau, J.-G.; Imbos, R.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2003**, *5*, 681–684. See also: Boiteau, J.-G.; Imbos, R.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2003**, *5*, 1385. (b) Boiteau, J.-G.; Minnaard, A. J.; Feringa, B. L. *J. Org. Chem.* **2003**, *68*, 9481–9484. (c) Duursma, A.; Hoen, R.; Schuppan, J.; Hulst, R.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2003**, *5*, 3111–3113.

(13) (a) Feringa, B. L. *Acc. Chem. Res.* **2000**, *33*, 346–353. (b) Naasz, R.; Arnold, L. A.; Minnaard, A. J.; Feringa, B. L. *Angew. Chem., Int. Ed.* **2001**, *40*, 927–930. (c) Arnold, L. A.; Naasz, R.; Minnaard, A. J.; Feringa, B. L. *J. Am. Chem. Soc.* **2003**, *125*, 3700–3701.


(14) Peña, D.; Lopez, F.; Harutyunyan, A. R.; Minnaard, A. J.; Feringa, B. L. *Chem. Commun.* **2004**, 1836–1837.

(15) (a) Sakai, M.; Hayashi, H.; Miyaura, N. *Organometallics* **1997**, *16*, 4229–4231. (b) Takaya, Y.; Ogasawara, M.; Hayashi, T.; Sakai, M.; Miyaura, M. *J. Am. Chem. Soc.* **1998**, *120*, 5579–5581. (c) For a recent review, see: Hayashi, T.; Yamasaki, K. *Chem. Rev.* **2003**, *103*, 2829–2844.

(16) Reetz, M. T.; Moulin, D.; Gosberg, A. *Org. Lett.* **2001**, *3*, 4083–4085.

(17) Kuriyama, M.; Nagai, K.; Yamada, K.; Miwa, Y.; Taga, T.; Tomioka, K. *J. Am. Chem. Soc.* **2002**, *124*, 8932–8939.

**Scheme 1**



We envisioned that introduction of aryl groups to **1** (Scheme 1), using the rhodium/phosphoramidite-catalyzed conjugate addition of arylboronic acids, could provide a pathway to 2-substituted 4-piperidones that is complementary to our work with dialkylzinc reagents. During our studies, Hayashi *et al.* reported the enantioselective addition of arylzinc chlorides to 2,3-dihydro-4-pyridones.<sup>7</sup> In that study, it was noted also that this type of substrate is less reactive toward 1,4-addition compared to other enones. The rhodium/BINAP-catalyzed conjugate addition of phenylboronic acid failed to proceed to full conversion, although the enantioselectivity was excellent.

Initial screening of our catalyst system on substrate **1** was performed under standard conditions in a mixture of dioxane/water (10/1) at 100 °C with a catalyst generated from 3 mol % Rh(acac)(C<sub>2</sub>H<sub>4</sub>)<sub>2</sub> and 7.5 mol % **L**. As in the report of Hayashi, with 3 equiv of phenylboronic acid, the reaction did not go to completion according to <sup>1</sup>H NMR (entry 1, Table 1). The enantioselectivity was, however, excellent

**Table 1.** Optimization of the Reaction Conditions for the Rhodium-Catalyzed Conjugate Addition to **1**

entry	"PhB" (equiv)	conditions <sup>a</sup>	conversion % <sup>b</sup>	ee (%) <sup>c</sup>
1	PhB(OH) <sub>2</sub> (3.0)	A	80	96
2	(PhBO) <sub>3</sub> (1.0)	B	60	99
3	(PhBO) <sub>3</sub> (3.0)	B	75	99
4	(PhBO) <sub>3</sub> (1.0)	C	84	99
5	(PhBO) <sub>3</sub> (2.0)	C	92	99
6	(PhBO) <sub>3</sub> (3.0)	C	100	99

<sup>a</sup> All reactions were performed on a 0.2 mmol scale with 3 mol % Rh(acac)(C<sub>2</sub>H<sub>4</sub>)<sub>2</sub> and 7.5 mol % (*R*)-**L** at 100 °C for 2 h. Conditions A: 0.55 mL of dioxane/H<sub>2</sub>O (10/1). Conditions B: 0.5 mL of dioxane, 1 equiv of H<sub>2</sub>O with respect to boron. Conditions C: 0.5 mL of dioxane, slow addition of water by syringe pump, 100 °C, 1 h. <sup>b</sup> Determined by <sup>1</sup>H NMR. <sup>c</sup> Determined by chiral HPLC.

(96% ee). We then decided to generate phenylboronic acid *in situ* from phenylboroxine ((PhBO)<sub>3</sub>)<sup>18</sup> and water (one

(18) Arylboroxines are obtained by dehydration of arylboronic acids by azeotropic removal of water from their xylene solution or heating at 300 °C *in vacuo*. For a review, see: Lappert, M. F. *Chem. Rev.* **1956**, *56*, 959–1064.

equivalent with respect to boron), providing mild reaction conditions (entries 2–5).<sup>19</sup> The use of this reagent did not improve the conversion but did improve the enantioselectivity to an excellent ee of 99%. Upon slow addition of water, thereby preventing premature hydrolysis of the boroxine, the reaction could be driven to 84% conversion using 1 equiv of boroxine (entry 4) and to full conversion with retention of 99% ee using 3 equiv of the reagent (entry 6).

To show the applicability of this reaction for synthesis on a laboratory scale, it was performed on a 0.5 g (2.2 mmol) scale. After flash chromatography, the product was isolated in 86% yield with 99% ee.

With these optimized conditions in hand, the scope of the asymmetric conjugate addition of arylboroxines to **1** was investigated. High ee values could be obtained with a variety of sterically and electronically diverse arylboroxines (entries 1–8, Table 2). *meta*- and *para*-tolyl groups can be introduced with high enantioselectivity and high yield (entries 3 and 4). A dramatic drop in enantioselectivity was observed for the more sterically demanding *ortho*-tolyl group (entry 2), illustrating a possible limitation of the catalytic method. Products with one or two electron-donating substituents on the aryl were obtained in high yield with high enantioselectivity (entries 5 and 6). However, electron-withdrawing groups such as chloride or fluoride slow the reaction, leading to incomplete conversions (entries 7 and 8). Despite this observation, the enantioselectivity is largely independent of the electronic properties of the substituents, and all *para*- or *meta*-substituted products are obtained with excellent ee values between 94 and 98%.

In summary, we have shown that conjugate addition of arylboroxines with a rhodium/phosphoramidite catalyst can be used to prepare 2-aryl-4-piperidones in high isolated yield (82–92%) and with excellent enantioselectivity (up to 99%). We are currently directing our efforts toward enhancing the

(19) Use of arylboroxines was previously reported to have a beneficial effect on both conversion and enantioselectivity in the conjugate addition to highly deactivated 1-alkenylphosphonates; see: Hayashi, T.; Senda, T.; Takaya, Y.; Ogasawara, M. *J. Am. Chem. Soc.* **1999**, *121*, 11591–11592.

**Table 2.** Scope of Arylboroxines in the Rhodium-Catalyzed Asymmetric 1,4-Addition to **1**<sup>a</sup>

entry	Ar	product	yield (%) <sup>b</sup>	ee (%) <sup>c</sup>
1	Ph	<b>2a</b>	86 <sup>d</sup>	99
2	<i>o</i> -MeC <sub>6</sub> H <sub>4</sub>	<b>2b</b>	82 <sup>d</sup>	24
3	<i>m</i> -MeC <sub>6</sub> H <sub>4</sub>	<b>2c</b>	92 <sup>d</sup>	98
4	<i>p</i> -MeC <sub>6</sub> H <sub>4</sub>	<b>2d</b>	86 <sup>d</sup>	95
5	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	<b>2e</b>	85 <sup>d</sup>	96
6	<i>m,p</i> -(MeO) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	<b>2f</b>	86 <sup>d</sup>	98
7	<i>p</i> -FC <sub>6</sub> H <sub>4</sub>	<b>2g</b>	71	94
8	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	<b>2h</b>	55	96

<sup>a</sup> All reactions were performed in duplicate with both enantiomers of the ligand on a 0.2 mmol scale with 3 mol % Rh(acac)(C<sub>2</sub>H<sub>4</sub>)<sub>2</sub> and 7.5 mol % **L** at 100 °C for 2 h. (*R*)-**L** gave the (*R*)-enantiomer of the product in all cases; see Supporting Information. <sup>b</sup> Isolated yield. <sup>c</sup> Determined by chiral HPLC. <sup>d</sup> Thin-layer chromatography shows a spot to spot conversion in 2 h.

scope and applications of this method in the synthesis of more complex heterocycles.

**Acknowledgment.** We thank Mrs. T. D. Tiemersma-Wegman for the technical support. Financial support from DSM, the Ministry of Economic Affairs, and NWO/CW, administered through the CW/Combichem program, is gratefully acknowledged.

**Supporting Information Available:** Experimental details and chromatographic and spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL050734M